

In the Specification:

Please enter substitute pages 45-96 of the application which comprise the unamended sequence listing for original pages 45-96.

In the claims:

1.(Amended) ~~An isolated nucleic acid encoding a protein which binds CD28 or CTLA4 comprising a contiguous nucleotide sequence derived from at least one T cell costimulatory molecule gene, the nucleotide sequence represented by a formula A-B-C-D-E, wherein~~

Sull
B1
a2

A comprises a nucleotide sequence of at least one first exon of a T cell costimulatory molecule gene, wherein the at least one first exon encodes a signal peptide domain,

B comprises a nucleotide sequence of at least one second exon of a T cell costimulatory molecule gene, wherein the at least one second exon encodes an immunoglobulin variable region-like domain,

C comprises a nucleotide sequence of at least one third exon of a T cell costimulatory molecule gene, wherein the at least one third exon encodes an immunoglobulin constant region-like domain,

D comprises a nucleotide sequence of at least one fourth exon of a T cell costimulatory molecule gene, wherein the at least one fourth exon encodes a transmembrane domain, and

E comprises a nucleotide sequence of at least one fifth exon of a T cell costimulatory molecule gene, wherein the at least one fifth exon encodes a cytoplasmic domain,

with the proviso that E [does not comprise a] is not any of the following nucleotide [sequence] sequences selected from a group consisting of SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29 and SEQ ID NO:31.

Sull
B3
a2

33. (Amended) ~~An isolated nucleic acid encoding a protein which binds CD28 or CTLA4 comprising a contiguous nucleotide sequence derived from at least one T cell~~

costimulatory molecule gene, the nucleotide sequence represented by a formula A-B-C-D-E, wherein

A comprises a nucleotide sequence of at least one first exon of a T cell costimulatory molecule gene, wherein the at least one first exon encodes a signal peptide domain,

B comprises a nucleotide sequence of at least one second exon of a T cell costimulatory molecule gene, wherein the at least one second exon encodes an immunoglobulin variable region-like domain,

C comprises a nucleotide sequence of at least one third exon of a T cell costimulatory molecule gene, wherein the at least one third exon encodes an immunoglobulin constant region-like domain,

D, which may or may not be present, comprises a nucleotide sequence of at least one fourth exon of a T cell costimulatory molecule gene, wherein the at least one fourth exon encodes a transmembrane domain, and

E, which may or may not be present, comprises a nucleotide sequence of at least one fifth exon of a T cell costimulatory molecule gene, wherein the at least one fifth exon encodes a cytoplasmic domain,

with the proviso that A [does not comprise a] is not any of the following nucleotide [sequence] sequences selected from a group consisting of SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39 and SEQ ID NO:41.

42. (Amended) [The] An isolated nucleic acid [of claim 40 which does not comprise a nucleotide sequence encoding the first signal peptide domain] which encodes a protein which binds to CD28 or CTLA4, wherein said nucleic acid molecule comprises a nucleotide sequence shown in SEQ ID NOs: 4 and 14.

Please add the following new claim:

77. (New) An isolated nucleic acid encoding a protein which binds CD28 or CTLA4 comprising a contiguous nucleotide sequence derived from at least one T cell costimulatory molecule gene, the nucleotide sequence represented by a formula A-B-C-D-E, wherein:

~~A comprises a nucleotide sequence of at least one first exon of a T cell costimulatory molecule gene, wherein the at least one first exon encodes a signal peptide domain,~~

~~B comprises a nucleotide sequence of at least one second exon of a T cell costimulatory molecule gene, wherein the at least one second exon encodes an immunoglobulin variable region-like domain,~~

~~C comprises a nucleotide sequence of at least one third exon of a T cell costimulatory molecule gene, wherein the at least one third exon encodes an immunoglobulin constant region-like domain,~~

~~D comprises a nucleotide sequence of at least one fourth exon of a T cell costimulatory molecule gene, wherein the at least one fourth exon encodes a transmembrane domain, and~~

~~E comprises a nucleotide sequence of at least one fifth exon of a T cell costimulatory molecule gene, wherein the at least one fifth exon encodes a cytoplasmic domain,~~

with the proviso that E is not any of the following nucleotide sequences selected from a group consisting of SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29 and SEQ ID NO:31; or wherein

AS cont
~~A comprises a nucleotide sequence of at least one first exon of a T cell costimulatory molecule gene, wherein the at least one first exon encodes a signal peptide domain,~~

~~B comprises a nucleotide sequence of at least one second exon of a T cell costimulatory molecule gene, wherein the at least one second exon encodes an immunoglobulin variable region-like domain,~~

~~C comprises a nucleotide sequence of at least one third exon of a T cell costimulatory molecule gene, wherein the at least one third exon encodes an immunoglobulin constant region-like domain,~~

~~D, which may or may not be present, comprises a nucleotide sequence of at least one fourth exon of a T cell costimulatory molecule gene, wherein the at least one fourth exon encodes a transmembrane domain, and~~

~~E, which may or may not be present, comprises a nucleotide sequence of at least one fifth exon of a T cell costimulatory molecule gene, wherein the at least one fifth exon encodes a cytoplasmic domain,~~